

University of Bergen, Department of Physics

Scientific/Technical Report No.1997-01

ISSN 0803-2696

hep-ph/9701245

January 1997

Probing the CP of the Higgs at an e^-e^- collider ¹

Odd Magne Ogreid

Department of Physics, University of Bergen, Allegaten 55, N-5007 BERGEN
NORWAY

Abstract

The sensitivity of various angular and energy distributions to the CP of a Higgs candidate are considered for the process $e^-e^- \rightarrow e^-e^-h$, where h is either a scalar or a pseudoscalar Higgs. This process may differ from the corresponding production at an e^+e^- collider in the availability of both beams being polarized. Beam polarization enhances the sensitivity of the azimuthal distribution.

1 Introduction

In planning for the Next Linear Collider [1, 2], it is important that one in addition to the electron-positron mode also considers the electron-electron mode [3]. One reason for this is that one can produce states not readily accessible in the annihilation channel. Another reason is that polarization of the electron beams will be readily available. The initial state consisting of two electrons is also a very clean state, there is no background from annihilation processes

When the Higgs particle is discovered and its mass has been measured, one will want to explore other properties of this particle. One of these is whether it is a scalar or a pseudoscalar. The Higgs predicted by the Standard Model is a scalar particle. However, certain extensions of the Standard Model allow for more than one Higgs particle. We can have both scalar and pseudoscalar Higgs particles in these models. This is the case in the MSSM [4]. When a possible candidate for the Higgs is observed, it is therefore of the utmost importance to determine its properties, and in particular its CP .

The possibility of disentangling the CP of the Higgs particle in the electron-electron mode will here be considered. This question has already been considered for the Bjorken process, where distributions in polar angle provide an easy resolution [5]. Also, azimuthal and energy distributions can be used for an independent determination of the CP [6].

The tree level diagrams for the process

$$e^-(p_1) + e^-(p_2) \rightarrow e^-(p'_1) + e^-(p'_2) + h(p_h)$$

are those where the Higgs particle (scalar or pseudoscalar) is produced via the fusion of two Z particles. There are two diagrams at the tree level because of identical electrons in the final state. For the ZZh coupling one can take [7]

$$i2^{5/4}\sqrt{G_F} \begin{cases} m_Z^2 g^{\mu\nu} & \text{for } h = H \text{ (} CP \text{ even),} \\ \eta \epsilon^{\mu\nu\rho\sigma} k_{1\rho} k_{2\sigma} & \text{for } h = A \text{ (} CP \text{ odd).} \end{cases} \quad (1)$$

Here, distributions which are obtained when all outgoing leptons are detected will be considered. The more favoured production via WW fusion will therefore not be discussed in this paper.

One expects the Higgs to be light. Therefore, in the numerical calculations, a Higgs mass of 135 GeV has been used. This value is consistent with a large number of models.

¹Presented at *QFTHEP '96*, XIth Workshop on Quantum Field Theory and High Energy Physics, St. Petersburg, Russia, September 12-18, 1996, to appear in the Proceedings.

2 Kinematics

The two electrons in the final state will be distinguished according to which has the higher energy, thus $E'_1 \geq E'_2$. The angles between the final state and the initial state electrons are denoted θ_1 and θ_2 ,

$$\begin{aligned}\mathbf{p}_1 \cdot \mathbf{p}'_1 &= |\mathbf{p}_1| |\mathbf{p}'_1| \cos \theta_1, \\ \mathbf{p}_2 \cdot \mathbf{p}'_2 &= |\mathbf{p}_2| |\mathbf{p}'_2| \cos \theta_2.\end{aligned}$$

Since the process contains three particles in the final state, the incoming and outgoing particles will not all lie in the same plane. But the two momenta $\mathbf{p}_1$ and $\mathbf{p}'_1$ will determine a plane, and similarly for $\mathbf{p}_2$ and $\mathbf{p}'_2$. The angle between these two planes will be denoted ϕ ,

$$\cos \phi = \frac{(\mathbf{p}_1 \times \mathbf{p}'_1) \cdot (\mathbf{p}_2 \times \mathbf{p}'_2)}{|\mathbf{p}_1 \times \mathbf{p}'_1| |\mathbf{p}_2 \times \mathbf{p}'_2|}.$$

The degree of longitudinal polarization for each of the initial electrons will be denoted P_i , thus $P_i = +1$ for pure states of positive helicity and $P_i = -1$ for pure states of negative helicity.




Figure 1: Azimuthal distributions for different values of the beam energy and various degrees of electron polarizations. The Higgs mass considered is 135 GeV. The solid lines correspond to a scalar Higgs and the dashed lines to a pseudoscalar one.

3 The cross section

The four-fold differential cross section

$$\frac{d^4\sigma}{d\epsilon d\phi d\cos\theta_1 d\cos\theta_2} \quad (2)$$

is calculated in both the scalar and the pseudoscalar case. The variable ϵ is the (positive) energy difference between the two final-state electrons, thus $\epsilon = E'_1 - E'_2$. This fully differential cross section is not easily studied experimentally, and therefore only cross sections obtained by integrating over two or three of the variables will be considered. In the extreme forward directions the electrons will be difficult to detect. Therefore, a cut on the polar angles θ_i is imposed in such a way that $|\cos \theta_i| \leq 0.9$, whenever an integration over these variables is performed.

3.1 Azimuthal distributions

The azimuthal distribution

$$\frac{2\pi}{\sigma_{\text{cut}}} \frac{d\sigma_{\text{cut}}}{d\phi} \quad (3)$$

has been determined by numerical integration, and is shown in Figure 1 for two different beam energies (150 GeV and 250 GeV) and at various polarizations for both scalar and pseudoscalar Higgs. For the unpolarized case the distributions are rather similar for a CP -odd and a CP -even Higgs at a beam energy of 150 GeV. At a higher energy, $E=250$ GeV, one can see a distinct difference between the two cases. This difference can be enhanced by turning on electron polarization. In the case of opposite polarizations (100%), one sees that the CP -even distribution behaves like $1 + \cos \phi$ while the CP -odd distribution behaves more like $1 - \cos 2\phi$. The term proportional to $\cos 2\phi$ has its origin in interference between transverse Z -particles with parallel and orthogonal linear polarizations, whereas a term proportional to $\cos \phi$ will arise from interference between scalar and transversely polarized vector bosons [8]. In the more realistic case of 85% electron polarization one can get nearly the same effect as by increasing the beam energy to around 250 GeV.

3.2 Energy distributions

From purely kinematical considerations one can show that

$$0 \leq \epsilon \leq \frac{1}{2}E - \frac{1}{8}\frac{m_h^2}{E} \equiv \epsilon_{\text{max}},$$

where m_h is the corresponding Higgs mass. A new variable, the “scaled energy difference”, $x = \epsilon/\epsilon_{\text{max}}$, ($0 \leq x \leq 1$), is therefore introduced. The x -distribution

$$\frac{1}{\sigma_{\text{cut}}} \frac{d\sigma_{\text{cut}}}{dx} \quad (4)$$

is shown in Figure 2. As can be seen from the plots, the distributions are rather similar in both the CP -even and the CP -odd case. A minor gain can be obtained from polarizing the beams, but probably too small to be useful in determining the CP of the Higgs.




Figure 2: The energy difference distribution shown at a beam energy of 150 GeV, and for two different cases of electron polarization. For both plots, the Higgs mass considered is 135 GeV. The solid lines correspond to a scalar Higgs and the dashed lines to a pseudoscalar one.

3.3 Polar angle distributions

Finally, the two-dimensional polar angle distribution

$$\frac{1}{\sigma_{\text{cut}}} \frac{d^2\sigma}{d\cos\theta_1 d\cos\theta_2}, \quad (5)$$

is considered. Representative plots are shown in Figure 3. The plots are made for the unpolarized case only. Little can be gained by using electron polarization when considering this distribution. The CP -odd distribution totally changes nature as one increases the beam energy from 100 GeV to 200 GeV. The CP -odd and CP -even distributions are easier to distinguish from each other at the lower beam energies. As the beam energy increases, they both peak in the forward region where $\cos\theta_1 = \cos\theta_2 = \pm 1$. But the production occurring in the forward region for $CP = -1$ is still an order of magnitude lower than what is the case for $CP = +1$.




Figure 3: Polar angle distributions for a scalar Higgs (left figures) and for a pseudoscalar Higgs (right figures) at various beam energies. For all plots, the Higgs mass considered is 135 GeV.

4 Summary

Among the three different distributions presented here, the azimuthal distribution and the polar angle distribution are those that contain the clearest signatures of the CP of the Higgs. The azimuthal distribution can be made quite distinct if beam polarization is available, in particular if the two beams are oppositely polarized. Also, one gets better separation at higher energies.

For the two-dimensional polar angle distribution, the difference between $CP = 1$ and $CP = -1$ is most striking at lower energies. In the CP -even case, the scattered electrons are forward peaked. As the beam energy increases, the distributions get more forward peaked. In the CP -odd case, there is a qualitative change from low to higher energies. At low energies (or heavy Higgs) the electrons tend to be scattered out to large angles, $|\cos\theta_i| \simeq 0$, whereas at higher energies the distributions peak *near* the forward direction.

It is a pleasure to thank my collaborators C. A. Bøe and Per Osland for helpful discussions and suggestions along the way. I will also thank I. F. Ginzburg for addressing the question of the origin of the different azimuthal distributions.

This work has been supported by the Research Council of Norway.

References

- [1] S.Kuhlman *et al.* (The NLC Accelerator Design Group and the NLC Physics Working Group) *Physics and Technology of the Next Linear Collider: A Report Submitted to Snowmass '96* by the NLC Zeroth-Order Design Group and the NLC Physics Working Group, BNL 52-502, FERMILAB-PUB-96/112, LBNL-PUB-5425, SLAC-Report-485, UCRL-ID-124160, UC-414 (June, 1996), (hep-ex/9605011).
- [2] ECFA/DESY Workshop 1996, report to appear.
- [3] Frank Cuypers, *Physics at an e^-e^- facility*, Talk given at 3rd workshop on physics and experiments with e^+e^- colliders, (LCWS95), Iwate, Japan, MPI-PHT-95-127 (hep-ph/9512201).
- [4] H.-P. Nilles, Phys. Rep. C110 (1984) 1;
H. E. Haber and G. L. Kane, Phys. Rep. C117 (1985) 75;
R. Barbieri, Riv. Nuovo Cimento 11 (1988) No. 4, p. 1;
For a recent review, see, e.g., R. Arnowitt and P. Nath, Lecture at Swieca School, Campos do Jordao, Brazil, 1993; in *Sao Paulo 1993*, Proceedings, Particles and fields, 3-63; CTP-TAMU-93-052 and NUB-TH-3073-93, (hep-ph/9309277).
- [5] V. Barger, K. Cheung, A. Djouadi, B. A. Kniehl and P. M. Zerwas, Phys. Rev. D49 (1994) 79.
- [6] A. Skjold and P. Osland, Nucl. Phys. B 453 (1995) 3.
- [7] D. Chang, W.-Y. Keung, I. Phillips, Phys. Rev. D48 (1993) 3225.
- [8] V. M. Budnev, I. F. Ginzburg, G. V. Meledin and V. G. Serbo, Phys. Reports 15 (1974) 181.